

# SOME PROPERTIES OF EXTENDED REMAINDER OF BINET'S FIRST FORMULA FOR LOGARITHM OF GAMMA FUNCTION

FENG QI AND BAI-NI GUO

ABSTRACT. In the paper, we extend Binet's first formula for the logarithm of the gamma function and investigate some properties, including inequalities, star-shaped and sub-additive properties and the complete monotonicity, of the extended remainder of Binet's first formula for the logarithm of the gamma function and related functions.

## 1. INTRODUCTION

1.1. For positive numbers  $x$  and  $y$  with  $y > x$ , let

$$g_{x,y}(t) = \int_x^y u^{t-1} du = \begin{cases} \frac{y^t - x^t}{t}, & t \neq 0, \\ \ln y - \ln x, & t = 0. \end{cases} \quad (1)$$

The reciprocal of  $g_{x,y}(t)$  can be rewritten as

$$\frac{1}{g_{e^a, e^b}(t)} = F_{a,b}(t) = \begin{cases} \frac{t}{e^{bt} - e^{at}}, & t \neq 0, \\ \frac{1}{b-a}, & t = 0, \end{cases}$$

where  $a$  and  $b$  are real numbers with  $b > a$ .

It is well-known [11, p. 11] that Binet's first formula of  $\ln \Gamma(x)$  for  $x > 0$  is given by

$$\ln \Gamma(x) = \left(x - \frac{1}{2}\right) \ln x - x + \ln \sqrt{2\pi} + \theta(x), \quad (2)$$

where

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$$

stands for Euler's gamma function and

$$\theta(x) = \int_0^\infty \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{e^{-xt}}{t} dt \quad (3)$$

for  $x > 0$  is called the remainder of Binet's first formula for the logarithm of the gamma function.

In [29, 30], some inequalities and completely monotonic properties of the function  $g_{x,y}(t)$  were established and applied to construct Steffensen pairs in [4], to refine Gautschi-Kershaw's inequalities in [24, 25], and to study monotonic properties, logarithmic convexities and Schur-convexities of extended mean values  $E(r, s; x, y)$  in [13, 14, 15, 16, 26, 31]. See also [17, 18] for related contents.

---

2000 *Mathematics Subject Classification.* 26A48, 26A51, 33B15.

*Key words and phrases.* inequality, extended remainder, Binet's first formula, gamma function, completely monotonic function, star-shaped function, sub-additive function.

The authors were supported in part by the China Scholarship Council and the Science Foundation of Tianjin Polytechnic University.

This manuscript is a slightly revised version of the formally-published paper [21].

In recent years, some inequalities and monotonic properties of the function

$$\frac{1}{t^2} - \frac{e^{-t}}{(1 - e^{-t})^2} \quad (4)$$

for  $t > 0$  and related ones were researched in [5, 10, 32] and related references therein. These results were used in [2, 8, 27, 28] to consider completely monotonic properties of remainders of Binet's first formula, the psi function and related ones.

Recently, logarithmic convexities of  $g_{x,y}(t)$  and  $F_{a,b}(t)$  were found in [7, 23]. By virtue of these conclusions, some simple and elegant proofs for the logarithmic convexities, Schur-convexities of extended mean values  $E(r, s; x, y)$  were simplified in [6, 19].

1.2. Now it is very natural to ask a question: Are there any relationship between the above studies? Direct computation yields

$$[g_{e^a, e^b}(t)]' = -[\ln F_{a,b}(t)]' = \frac{be^{bt} - ae^{at}}{e^{bt} - e^{at}} - \frac{1}{t} = \frac{b-a}{e^{(b-a)t} - 1} - \frac{1}{t} + b \triangleq \delta_{a,b}(t) \quad (5)$$

for  $t \neq 0$ . Therefore, if taking  $a = -\frac{1}{2}$  and  $b = \frac{1}{2}$ , then  $\delta_{-1/2, 1/2}(t)$  for  $t > 0$  equals the integrand in the remainder of Binet's first formula for the logarithm of the gamma function and the first order derivative of  $\delta_{-1/2, 1/2}(t)$  for  $t > 0$  also equals the function (4). These relationships connect closely the above three seemingly unrelated problems.

If replacing  $\delta_{-1/2, 1/2}(t)$  by  $\delta_{a,b}(t)$  on  $t \in (0, \infty)$  for  $b > a$  in (3), then a more problem emerges: How to calculate the improper integral

$$\int_0^\infty \left[ \frac{b-a}{e^{(b-a)t} - 1} - \frac{1}{t} + b \right] \frac{e^{-tx}}{t} dt \quad (6)$$

for  $b > a$ ? The following Theorem 1 answers this question affirmatively.

**Theorem 1.** *Let  $b > a$ ,  $\alpha > 0$  and  $x \in \mathbb{R}$  be real numbers. Then the improper integral (6) converges if and only if  $a + b = 0$  and*

$$\int_0^\infty \left( \frac{\alpha}{e^{\alpha t} - 1} - \frac{1}{t} + \frac{\alpha}{2} \right) \frac{e^{-tx}}{t} dt = \alpha \ln \Gamma\left(\frac{x}{\alpha}\right) - \left(x - \frac{\alpha}{2}\right) \ln \frac{x}{\alpha} + x - \frac{\alpha}{2} \ln(2\pi). \quad (7)$$

*Remark 1.* It is easy to see that the formula (2) is the special case  $\alpha = 1$  of (7). So we call (7) the extended remainder of Binet's first formula for the logarithm of the gamma function  $\Gamma$ .

1.3. It is well-known [8, 20, 22] that a function  $f$  is said to be completely monotonic on an interval  $I$  if  $f$  has derivatives of all orders on  $I$  and

$$(-1)^n f^{(n)}(x) \geq 0$$

for  $x \in I$  and  $n \geq 0$ , that a function  $f(x)$  is said to be star-shaped on  $(0, \infty)$  if

$$f(\alpha x) \leq \alpha f(x)$$

for  $x \in (0, \infty)$  and  $0 < \alpha < 1$ , that a function  $f$  is said to be super-additive on  $(0, \infty)$  if

$$f(x+y) \geq f(x) + f(y)$$

for all  $x, y > 0$ , and that a function  $f$  is said to be sub-additive if  $-f$  is super-additive.

The function  $\delta_{a,b}(t)$  defined by (5) has the following properties.

**Theorem 2.** *Let  $a$  and  $b$  be real numbers with  $a \neq b$ , and let  $0 < \tau < 1$ .*

- (1) *The function  $\delta_{a,b}(t)$  is increasingly concave on  $(0, \infty)$  and convex on  $(-\infty, 0)$ ;*

(2) *The inequality*

$$\delta_{a,b}(\tau t) < \delta_{a,b}(t) \quad (8)$$

*is valid on  $(0, \infty)$ ; if either  $a + b \neq 0$  or  $ab \neq 0$ , then the inequality (8) is sharp;*

(3) *If  $a + b \geq 0$ , then*

$$\tau \delta_{a,b}(t) < \delta_{a,b}(\tau t) \quad (9)$$

*is valid on  $(0, \infty)$ , i.e., the function  $-\delta_{a,b}(t)$  is star-shaped; if  $\max\{a, b\} \leq 0$ , then the inequality (9) is reversed, i.e., the function  $\delta_{a,b}(t)$  is star-shaped; if  $a + b = 0$ , then the inequality (9) is sharp.*

*Remark 2.* Some properties of special cases of the function  $\delta_{a,b}(t)$  and related ones have been investigated and applied extensively in [5, 6, 7, 8, 10, 23, 29, 30, 32] and related references therein.

1.4. If denoting the extended remainder of Binet's first formula for the logarithm of the gamma function  $\Gamma$  by

$$\theta_\alpha(x) = \int_0^\infty \left( \frac{\alpha}{e^{\alpha t} - 1} - \frac{1}{t} + \frac{\alpha}{2} \right) \frac{e^{-tx}}{t} dt \quad (10)$$

for  $\alpha > 0$  and  $x > 0$ , then formula (7) in Theorem 1 can be simplified as

$$\theta_\alpha(x) = \alpha \theta\left(\frac{x}{\alpha}\right) \quad \text{or} \quad \theta_\alpha(\alpha x) = \alpha \theta(x).$$

This motivates us to study properties of  $\theta_\alpha(x)$  and the function

$$f_{p,q;\alpha}(x) = \theta_\alpha(px) - q\theta_\alpha(x)$$

on  $(0, \infty)$ , where  $p > 0$ ,  $\alpha > 0$  and  $q \in \mathbb{R}$ , which may be concluded as the following theorem.

**Theorem 3.** *The extended remainder  $\theta_\alpha(x)$  of Binet's first formula for the logarithm of the gamma function  $\Gamma$  satisfies*

$$\frac{(-1)^k}{(1+\lambda)^k} \theta_\alpha^{(k)}\left(\frac{x}{1+\lambda}\right) > \frac{(-1)^k}{2} \left[ \frac{1}{\lambda^k} \theta_\alpha^{(k)}\left(\frac{x}{\lambda}\right) + \theta_\alpha^{(k)}(x) \right] \quad (11)$$

for  $x > 0$ ,  $\lambda > 0$  with  $\lambda \neq 1$ ,  $k \geq 0$  and  $\alpha > 0$ .

*The function  $f_{p,q;\alpha}(x)$  is completely monotonic on  $(0, \infty)$  if either  $0 < p \leq 1$  and  $q \leq 1$  or  $p > 1$  and  $q \leq \frac{1}{p}$ ; the function  $-f_{p,q;\alpha}(x)$  is completely monotonic on  $(0, \infty)$  if  $p \geq 1$  and  $q \geq 1$ .*

*The function  $-\theta_\alpha(x)$  is star-shaped and  $\theta_\alpha(x)$  is sub-additive.*

*Remark 3.* If taking  $a = -\frac{1}{2}$  and  $b = \frac{1}{2}$ , then the results obtained in [2, 3, 8, 27, 28] can be deduced directly from Theorem 3.

## 2. PROOFS OF THEOREMS

*Proof of Theorem 1.* By transformations of integral variables, it easily follows that

$$\begin{aligned} & \int_\varepsilon^\infty \left[ \frac{b-a}{e^{(b-a)t} - 1} - \frac{1}{t} + b \right] \frac{e^{-tx}}{t} dt \\ &= \int_{(b-a)\varepsilon}^\infty \left( \frac{b-a}{e^u - 1} - \frac{b-a}{u} + b \right) \frac{e^{-ux/(b-a)}}{u} du \\ &= (b-a) \int_{(b-a)\varepsilon}^\infty \left( \frac{1}{e^u - 1} - \frac{1}{u} + \frac{1}{2} \right) \frac{e^{-ux/(b-a)}}{u} du \\ & \quad + \frac{a+b}{2} \int_\varepsilon^\infty \frac{e^{-ux/(b-a)}}{u} du \end{aligned}$$

$$\begin{aligned}
&= (b-a) \int_{(b-a)\varepsilon}^{\infty} \left( \frac{1}{e^u - 1} - \frac{1}{u} + \frac{1}{2} \right) \frac{e^{-ux/(b-a)}}{u} du \\
&\quad + \frac{a+b}{2} \int_{(b-a)\varepsilon}^{\infty} \frac{e^{-ux}}{u} du,
\end{aligned} \tag{12}$$

where  $\varepsilon > 0$ .

By virtue of Binet's first formula for  $\ln \Gamma(z)$  in (3), the integral in the first term of (12) may be calculated as

$$\begin{aligned}
&\lim_{\varepsilon \rightarrow 0^+} \int_{(b-a)\varepsilon}^{\infty} \left( \frac{1}{e^u - 1} - \frac{1}{u} + \frac{1}{2} \right) \frac{e^{-ux/(b-a)}}{u} du \\
&= \int_0^{\infty} \left( \frac{1}{e^u - 1} - \frac{1}{u} + \frac{1}{2} \right) \frac{e^{-ux/(b-a)}}{u} du \\
&= \ln \Gamma \left( \frac{x}{b-a} \right) - \left( \frac{x}{b-a} - \frac{1}{2} \right) \ln \frac{x}{b-a} + \frac{x}{b-a} - \frac{1}{2} \ln(2\pi).
\end{aligned} \tag{13}$$

Furthermore, the second integral in (12) satisfies

$$\lim_{\varepsilon \rightarrow 0^+} \int_{(b-a)\varepsilon}^{\infty} \frac{e^{-ux}}{u} du = \lim_{\varepsilon \rightarrow 0^+} \int_{(b-a)x\varepsilon}^{\infty} t^{-1} e^{-t} dt = \int_0^{\infty} t^{-1} e^{-t} dt$$

which is divergent. As a result, the improper integral (6) is convergent if and only if  $a+b=0$ .

Taking  $a = -b$  in (6) and (13) and simplifying yields formula (7). The proof of Theorem 1 is complete.  $\square$

*Proof of Theorem 2.* Straightforward computation gives

$$\begin{aligned}
\delta'_{a,b}(t) &= \frac{1}{t^2} - \frac{(a-b)^2 e^{(a+b)t}}{(e^{at} - e^{bt})^2}, \\
\delta''_{a,b}(t) &= \frac{(a-b)^3 e^{(a+b)t} (e^{at} + e^{bt})}{(e^{at} - e^{bt})^3} - \frac{2}{t^3} \\
&= \frac{2e^{3(a+b)t/2}}{t^3} \left( \frac{at - bt}{e^{at} - e^{bt}} \right)^3 \left\{ \frac{e^{(a-b)t/2} + e^{(b-a)t/2}}{2} - \left[ \frac{e^{(a-b)t/2} - e^{(b-a)t/2}}{(a-b)t} \right]^3 \right\} \\
&\triangleq \frac{2e^{3(a+b)t/2}}{t^3} \left( \frac{at - bt}{e^{at} - e^{bt}} \right)^3 Q \left( \frac{a-b}{2} t \right).
\end{aligned}$$

Lazarević's inequality in [1, p. 131] and [9, p. 300] tells us that

$$Q(t) = \frac{e^{-t} + e^t}{2} - \left( \frac{e^t - e^{-t}}{2t} \right)^3 = \cosh t - \left( \frac{\sinh t}{t} \right)^3 < 0$$

for  $t \in \mathbb{R}$  with  $t \neq 0$ . Hence  $\delta''_{a,b}(t) < 0$  on  $(0, \infty)$  and  $\delta''_{a,b}(t) > 0$  on  $(-\infty, 0)$ . The convexity and concavity of  $\delta_{a,b}(t)$  are proved.

Since  $\delta''_{a,b}(t) < 0$  on  $(0, \infty)$ , the derivative  $\delta'_{a,b}(t)$  is decreasing on  $(0, \infty)$  for all real numbers  $a$  and  $b$  with  $a \neq b$ . Since

$$\delta'_{a,b}(t) = \frac{1}{t^2} - \frac{(a-b)^2 e^{(b-a)t}}{[1 - e^{(b-a)t}]^2} = \frac{1}{t^2} - \frac{(a-b)^2 e^{(a-b)t}}{[1 - e^{(a-b)t}]^2},$$

it is easy to obtain that

$$\lim_{t \rightarrow \infty} \delta'_{a,b}(t) = 0.$$

Consequently, the function  $\delta'_{a,b}(t)$  is positive, and so  $\delta_{a,b}(t)$  is increasing, on  $(0, \infty)$ . This means that inequality (8) holds for  $0 < \tau < 1$  and  $t > 0$ .

From

$$\delta_{a,b}(t) = \frac{be^{bt} - ae^{at}}{e^{bt} - e^{at}} - \frac{1}{t} = \frac{be^{(b-a)t} - a}{e^{(b-a)t} - 1} - \frac{1}{t} = \frac{b - ae^{(a-b)t}}{1 - e^{(a-b)t}} - \frac{1}{t},$$

it follows easily that

$$\lim_{t \rightarrow \infty} \delta_{a,b}(t) = \max\{a, b\}.$$

L'Hôpital's rule gives

$$\begin{aligned} \lim_{t \rightarrow 0^+} \delta_{a,b}(t) &= \lim_{t \rightarrow 0^+} \frac{t(be^{bt} - ae^{at}) - e^{bt} + e^{at}}{t(e^{bt} - e^{at})} \\ &= \lim_{t \rightarrow 0^+} \frac{b^2 e^{bt} - a^2 e^{at}}{(be^{bt} - ae^{at}) + (e^{bt} - e^{at})/t} \\ &= \frac{a+b}{2}. \end{aligned}$$

For  $0 < \tau < 1$ , let

$$h_{a,b}(t) = \delta_{a,b}(\tau t) - \tau \delta_{a,b}(t)$$

for  $t > 0$ . It is obvious that

$$\lim_{t \rightarrow 0^+} h_{a,b}(t) = \frac{(1-\tau)(a+b)}{2} \quad \text{and} \quad \lim_{t \rightarrow \infty} h_{a,b}(t) = (1-\tau) \max\{a, b\}.$$

Since  $\delta'_{a,b}(t)$  is decreasing, then

$$h'_{a,b}(t) = \tau [\delta'_{a,b}(\tau t) - \delta'_{a,b}(t)] > 0,$$

and so  $h_{a,b}(t)$  is strictly increasing. If  $a+b \geq 0$ , then  $h_{a,b}(t) > 0$  on  $(0, \infty)$  and inequality (9) is valid. If  $\max\{a, b\} \leq 0$ , then inequality (9) is reversed.

It is apparent that

$$\lim_{t \rightarrow \infty} \frac{\delta_{a,b}(\tau t)}{\delta_{a,b}(t)} = 1$$

if  $ab \neq 0$ , which implies that inequality (8) is sharp. If  $a+b \neq 0$ , then it is clear that

$$\lim_{t \rightarrow 0^+} \frac{\delta_{a,b}(\tau t)}{\delta_{a,b}(t)} = 1,$$

which also implies that inequality (8) is sharp.

By L'Hôpital's rule, it is not difficult to obtain that

$$\lim_{t \rightarrow 0^+} \delta'_{a,b}(t) = \frac{(a-b)^2}{12}.$$

If  $a+b=0$  and  $a \neq b$ , then

$$\lim_{t \rightarrow 0^+} \frac{\delta_{a,b}(\tau t)}{\delta_{a,b}(t)} = \lim_{t \rightarrow 0^+} \frac{\tau \delta'_{a,b}(\tau t)}{\delta'_{a,b}(t)} = \tau,$$

which means that inequality (9) is sharp. The proof of Theorem 2 is complete.  $\square$

*Proof of Theorem 3.* From the concavity of  $\delta_{a,b}(t)$  on  $(0, \infty)$ , it follows that

$$\delta_{-\alpha/2, \alpha/2}\left(\frac{(1+\lambda)t}{2}\right) > \frac{\delta_{-\alpha/2, \alpha/2}(\lambda t) + \delta_{-\alpha/2, \alpha/2}(t)}{2}$$

for  $t > 0$  and positive numbers  $\alpha$  and  $\lambda \neq 1$ . Multiplying by the factor  $t^{k-1}e^{-tx}$  for any nonnegative integer  $k \geq 0$  and integrating from 0 to  $\infty$  on both sides of the above inequality yields

$$\int_0^\infty \left[ \frac{\alpha}{e^{\alpha(1+\lambda)t/2} - 1} - \frac{2}{(1+\lambda)t} + \frac{\alpha}{2} \right] t^{k-1} e^{-tx} dt$$

$$\begin{aligned}
&> \frac{1}{2} \left[ \int_0^\infty \left( \frac{\alpha}{e^{\alpha\lambda t} - 1} - \frac{1}{\lambda t} + \frac{\alpha}{2} \right) t^{k-1} e^{-tx} dt \right. \\
&\quad \left. + \int_0^\infty \left( \frac{\alpha}{e^{\alpha t} - 1} - \frac{1}{t} + \frac{\alpha}{2} \right) t^{k-1} e^{-tx} dt \right]
\end{aligned}$$

which can be rewritten by transformations of integral variables as

$$\begin{aligned}
&\int_0^\infty \left( \frac{\alpha}{e^{\alpha u} - 1} - \frac{1}{u} + \frac{\alpha}{2} \right) \frac{u^k}{(1+\lambda)^k} \cdot \frac{e^{-xu/(1+\lambda)}}{u} du \\
&> \frac{1}{2} \left[ \int_0^\infty \left( \frac{\alpha}{e^{\alpha u} - 1} - \frac{1}{u} + \frac{\alpha}{2} \right) \frac{u^k}{\lambda^k} \cdot \frac{e^{-xu/\lambda}}{u} du \right. \\
&\quad \left. + \int_0^\infty \left( \frac{\alpha}{e^{\alpha t} - 1} - \frac{1}{t} + \frac{\alpha}{2} \right) t^k \frac{e^{-tx}}{t} dt \right].
\end{aligned}$$

Substituting formula (10) and its derivatives into the above inequalities leads to

$$(-1)^k \left[ \theta_\alpha \left( \frac{x}{1+\lambda} \right) \right]^{(k)} > \frac{1}{2} \left\{ (-1)^k \left[ \theta_\alpha \left( \frac{x}{\lambda} \right) \right]^{(k)} + (-1)^k \theta_\alpha^{(k)}(x) \right\}.$$

As a result, inequalities in (11) follow.

Easy calculation yields

$$\begin{aligned}
f_{p,q;\alpha}(x) &= \int_0^\infty \delta_{-\alpha/2,\alpha/2}(t) \frac{e^{-pxt}}{t} dt - q \int_0^\infty \delta_{-\alpha/2,\alpha/2}(t) \frac{e^{-xt}}{t} dt \\
&= \int_0^\infty \delta_{-\alpha/2,\alpha/2} \left( \frac{t}{p} \right) \frac{e^{-xt}}{t} dt - q \int_0^\infty \delta_{-\alpha/2,\alpha/2}(t) \frac{e^{-xt}}{t} dt \\
&= \int_0^\infty \left[ \delta_{-\alpha/2,\alpha/2} \left( \frac{t}{p} \right) - q \delta_{-\alpha/2,\alpha/2}(t) \right] \frac{e^{-xt}}{t} dt \\
&\triangleq \int_0^\infty h_{p,q;\alpha}(t) \frac{e^{-xt}}{t} dt.
\end{aligned}$$

By virtue of properties of  $\delta_{a,b}(t)$  obtained in Theorem 2, it follows by standard arguments that

- (1)  $h_{p,q;\alpha}(t) \geq 0$  if either  $0 < p \leq 1$  and  $q \leq 1$ , or  $p > 1$  and  $q \leq \frac{1}{p}$ , or  $0 < q < 1$  and  $q \leq \frac{1}{p}$ ;
- (2)  $h_{p,q;\alpha}(t) \leq 0$  if  $p \geq 1$  and  $q \geq 1$ .

It is clear that

- (1) if  $h_{p,q;\alpha}(t) \geq 0$  then  $f_{p,q;\alpha}(x)$  is completely monotonic on  $(0, \infty)$ ;
- (2) if  $h_{p,q;\alpha}(t) \leq 0$  then  $-f_{p,q;\alpha}(x)$  is completely monotonic on  $(0, \infty)$ .

As a result, the completely monotonic properties of  $f_{p,q;\alpha}(x)$  is proved.

It is easy to see that the star-shaped properties of the function  $\theta_\alpha(x)$  follow from those of the function  $\delta_{-\alpha/2,\alpha/2}(t)$  and formula (10).

In [12, p. 453], it was presented that a star-shaped function must be super-additive, therefore the function  $\theta_\alpha(x)$  is also sub-additive. The proof of Theorem 3 is complete.  $\square$

*Remark 4.* Dividing both sides of (7) by  $\alpha > 0$  gives

$$\int_0^\infty \left( \frac{1}{e^{\alpha t} - 1} - \frac{1}{\alpha t} + \frac{1}{2} \right) \frac{e^{-tx}}{t} dt = \ln \Gamma \left( \frac{x}{\alpha} \right) - \left( \frac{x}{\alpha} - \frac{1}{2} \right) \ln \frac{x}{\alpha} + \frac{x}{\alpha} - \frac{1}{2} \ln(2\pi). \quad (14)$$

Further taking  $x = \alpha y$  in (14) yields

$$\int_0^\infty \left( \frac{1}{e^{\alpha t} - 1} - \frac{1}{\alpha t} + \frac{1}{2} \right) \frac{e^{-\alpha ty}}{t} dt = \ln \Gamma(y) - \left( y - \frac{1}{2} \right) \ln y + y - \frac{1}{2} \ln(2\pi). \quad (15)$$

Since

$$\int_0^\infty \left( \frac{1}{e^{\alpha t} - 1} - \frac{1}{\alpha t} + \frac{1}{2} \right) \frac{e^{-\alpha t y}}{t} dt = \int_0^\infty \left( \frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{e^{-t y}}{t} dt, \quad (16)$$

so the identity (7) is essentially equivalent to (2).

## REFERENCES

- [1] P. S. Bullen, *A Dictionary of Inequalities*, Pitman Monographs and Surveys in Pure and Applied Mathematics **97**, Addison Wesley Longman Limited, 1998.
- [2] Ch.-P. Chen and F. Qi, *Completely monotonic functions related to the gamma functions*, RGMIA Res. Rep. Coll. **8** (2005), no. 2, Art. 3, 195–200; Available online at <http://rgmia.org/v8n2.php>.
- [3] Ch.-P. Chen, F. Qi and H. M. Srivastava, *Some properties of functions related to the gamma and psi functions*, Integral Transforms Spec. Funct. **21** (2010), no. 2, 153–164; Available online at <http://dx.doi.org/10.1080/10652460903064216>.
- [4] H. Gauchman, *Steffensen pairs and associated inequalities*, J. Inequal. Appl. **5** (2000), no. 1, 53–61.
- [5] B.-N. Guo, A.-Q. Liu and F. Qi, *Monotonicity and logarithmic convexity of three functions involving exponential function*, J. Korea Soc. Math. Educ. Ser. B Pure Appl. Math. **15** (2008), no. 4, 387–392.
- [6] B.-N. Guo and F. Qi, *A simple proof of logarithmic convexity of extended mean values*, Numer. Algorithms **52** (2009), 89–92; Available online at <http://dx.doi.org/10.1007/s11075-008-9259-7>.
- [7] B.-N. Guo and F. Qi, *The function  $(b^x - a^x)/x$ : Ratio's properties*, Available online at <http://arxiv.org/abs/0904.1115>.
- [8] S. Guo and F. Qi, *A class of completely monotonic functions related to the remainder of Binet's formula with applications*, Tamsui Oxf. J. Math. Sci. **25** (2009), no. 1, 9–14.
- [9] J.-Ch. Kuang, *Chángyòng Bùdēngshì (Applied Inequalities)*, 3rd ed., Shandong Science and Technology Press, Ji'nan City, Shandong Province, China, 2004. (Chinese)
- [10] A.-Q. Liu, G.-F. Li, B.-N. Guo and F. Qi, *Monotonicity and logarithmic concavity of two functions involving exponential function*, Internat. J. Math. Ed. Sci. Tech. **39** (2008), no. 5, 686–691.
- [11] W. Magnus, F. Oberhettinger and R. P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, Springer, Berlin, 1966.
- [12] A. W. Marshall and I. Olkin, *Inequalities: Theory of Majorization and its Applications*, Academic Press, New York, 1979.
- [13] F. Qi, *A note on Schur-convexity of extended mean values*, Rocky Mountain J. Math. **35** (2005), no. 5, 1787–1793.
- [14] F. Qi, *Logarithmic convexities of the extended mean values*, RGMIA Res. Rep. Coll. **2** (1999), no. 5, Art. 5, 643–652; Available online at <http://rgmia.org/v2n5.php>.
- [15] F. Qi, *Logarithmic convexity of extended mean values*, Proc. Amer. Math. Soc. **130** (2002), no. 6, 1787–1796.
- [16] F. Qi, *Schur-convexity of the extended mean values*, RGMIA Res. Rep. Coll. **4** (2001), no. 4, Art. 4, 529–533; Available online at <http://rgmia.org/v4n4.php>.
- [17] F. Qi, *The extended mean values: Definition, properties, monotonicities, comparison, convexities, generalizations, and applications*, Cubo Mat. Educ. **5** (2003), no. 3, 63–90.
- [18] F. Qi, *The extended mean values: Definition, properties, monotonicities, comparison, convexities, generalizations, and applications*, RGMIA Res. Rep. Coll. **5** (2002), no. 1, Art. 5, 57–80; Available online at <http://rgmia.org/v5n1.php>.
- [19] F. Qi, P. Cerone, S. S. Dragomir and H. M. Srivastava, *Alternative proofs for monotonic and logarithmically convex properties of one-parameter mean values*, Appl. Math. Comput. **208** (2009), no. 1, 129–133; Available online at <http://dx.doi.org/10.1016/j.amc.2008.11.023>.
- [20] F. Qi and B.-N. Guo, *A note on additivity of polygamma functions*, Available online at <http://arxiv.org/abs/0903.0888>.
- [21] F. Qi and B.-N. Guo, *Some properties of extended remainder of Binet's first formula for logarithm of gamma function*, Math. Slovaca **60** (2010), no. 4, 461–470; Available online at <http://dx.doi.org/10.2478/s12175-010-0025-7>.
- [22] F. Qi and B.-N. Guo, *Subadditive and superadditive properties of polygamma functions*, RGMIA Res. Rep. Coll. **10** (2007), Suppl., Art. 4; Available online at [http://rgmia.org/v10\(E\).php](http://rgmia.org/v10(E).php).
- [23] F. Qi and B.-N. Guo, *The function  $(b^x - a^x)/x$ : Logarithmic convexity*, RGMIA Res. Rep. Coll. **11** (2008), no. 1, Art. 5; Available online at <http://rgmia.org/v11n1.php>.

- [24] F. Qi, B.-N. Guo and Ch.-P. Chen, *The best bounds in Gautschi-Kershaw inequalities*, Math. Inequal. Appl. **9** (2006), no. 3, 427–436.
- [25] F. Qi, B.-N. Guo and Ch.-P. Chen, *The best bounds in Gautschi-Kershaw inequalities*, RGMIA Res. Rep. Coll. **8** (2005), no. 2, Art. 17, 311–320; Available online at <http://rgmia.org/v8n2.php>.
- [26] F. Qi and Q.-M. Luo, *A simple proof of monotonicity for extended mean values*, J. Math. Anal. Appl. **224** (1998), no. 2, 356–359.
- [27] F. Qi, D.-W. Niu and B.-N. Guo, *Monotonic properties of differences for remainders of psi function*, Internat. J. Pure Appl. Math. Sci. **4** (2007), no. 1, 59–66.
- [28] F. Qi, D.-W. Niu and B.-N. Guo, *Monotonic properties of differences for remainders of psi function*, RGMIA Res. Rep. Coll. **8** (2005), no. 4, Art. 16, 683–690; Available online at <http://rgmia.org/v8n4.php>.
- [29] F. Qi and S.-L. Xu, *Refinements and extensions of an inequality, II*, J. Math. Anal. Appl. **211** (1997), no. 2, 616–620.
- [30] F. Qi and S.-L. Xu, *The function  $(b^x - a^x)/x$ : Inequalities and properties*, Proc. Amer. Math. Soc. **126** (1998), no. 11, 3355–3359.
- [31] F. Qi, S.-L. Xu and L. Debnath, *A new proof of monotonicity for extended mean values*, Internat. J. Math. Math. Sci. **22** (1999), no. 2, 417–421.
- [32] Sh.-Q. Zhang, B.-N. Guo and F. Qi, *A concise proof for properties of three functions involving the exponential function*, Appl. Math. E-Notes **9** (2009), 177–183.

(F. Qi) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN CITY, 300160, CHINA

*E-mail address:* [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)

*URL:* <http://qifeng618.spaces.live.com>

(B.-N. Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

*E-mail address:* [bai.ni.guo@gmail.com](mailto:bai.ni.guo@gmail.com), [bai.ni.guo@hotmail.com](mailto:bai.ni.guo@hotmail.com)

*URL:* <http://guobaini.spaces.live.com>